


Uncertainty in Operational Atmospheric Analyses

Mean u250 variance ECMWF NOGAPS METFRANCE GEOS5



Objectives

- 1. Quantify the uncertainty (differences) in current operational analyses of the atmosphere – height, temperature, winds**
- 2. Consider implications of analysis uncertainty for NWP and plans for the future global observing network**

Analysis differences are a proxy for actual analysis error, which cannot be precisely quantified

Significance of Analysis Uncertainty/Error

- **Quality of NWP forecasts from short to medium-range**
- **Extended-range NWP?**
- **Short-range climate forecasts?**
- **Quality of forecast verification**
- **Accuracy of climate monitoring**

Causes of Analysis Differences and Error

- **Gaps/deficiencies in global observing network**
- **Errors /bias in observation data**
- **Choices in observation selection**
- **Observation quality control decisions**
- **Different and imperfect data assimilation techniques**
- **Errors in background forecast**

Methodology

- **Use multi-year, multi-model archive of operational analyses and forecasts, developed at NRL for research and diagnostic studies**
- **Quantify and examine differences in atmospheric analyses, trends over time ...**
- **Examine systematic (monthly/seasonal) patterns**

Surprisingly sparse literature on the topic of atmospheric analysis uncertainty and error

[Scholarly articles for uncertainty in atmospheric analyses](#)

[... of analysis uncertainty upon regional atmospheric ...](#) - Wang - Cited by 60

[Uncertainty analysis of climate change and policy ...](#) - Webster - Cited by 195

[On the assessment and uncertainty of atmospheric ...](#) - Abrams - Cited by 42

[PDF] [Uncertainty in Atmospheric CO₂ Concentrations from a Paramet...](#)

globalchange.mit.edu/files/document/MITJPSPGC_Rpt39.pdf

File Format: PDF/Adobe Acrobat - [Quick View](#)

Parametric **Uncertainty Analysis** of a Global Ocean Carbon Cycle Model. Gary Louis Holian. Submitted to the Department of Earth, **Atmospheric**, and Planetary ...

[Uncertainty in atmospheric CO₂ predictions from a parametric ...](#)

dspace.mit.edu/handle/1721.1/3565

by GL Holian - 2001 - Cited by 8 - [Related articles](#)

Uncertainty in atmospheric CO₂ predictions from a parametric uncertainty analysis of a global carbon cycle model. Show full item record. Citable URI: ...

[Quantitative uncertainty analyses of ancient atmospheric CO₂ ...](#)

ajsonline.org/content/309/9/775.abstract

by DJ Beerling - 2009 - Cited by 9 - [Related articles](#)

Quantitative **uncertainty analyses** of ancient **atmospheric CO₂** estimates from fossil leaves. David J. Beerling*,†,; Andrew Fox* and; Clive W. Anderson** ...

[Uncertainty in atmospheric temperature analyses | Langland | Tellus A](#)

journals.sfu.ca/coaction/index.php/tellusa/article/view/15390

by RH Langland - 2008 - Cited by 9 - [Related articles](#)

This report illustrates and quantifies the unanticipated large **uncertainty** and differences in tropospheric temperature **analyses** within current global operational ...

[PDF] [estimates, uncertainty analysis, and sensitivity analysis - ACP](#)

www.atmos-chem-phys.net/11/2625/2011/acp-11-2625-2011.pdf

File Format: PDF/Adobe Acrobat - [Quick View](#)

by IMD Rosa - 2011 - [Related articles](#)

Atmospheric. Chemistry and Physics. **Atmospheric** emissions from vegetation fires in. Portugal (1990–2008): estimates, **uncertainty analysis**, and sensitivity ...

[Uncertainty in atmospheric temperature analyses - LANGLAND ...](#)

onlinelibrary.wiley.com/.../Journal Home/Vol 60 Issue 4

by RH LANGLAND - 2008 - Cited by 9 - [Related articles](#)

Jul 8, 2008 – **Uncertainty in atmospheric temperature analyses**. ROLF H. LANGLAND1,*,; RYAN N. MAUE2,; CRAIG H. BISHOP1. Article first published ...

“Some aspects of the improvement in skill of numerical weather prediction, 2002: A.J. Simmons and A. Hollingsworth, QJRMS.

Langland, Maue,
Bishop, 2008: Tellus

“Analysis differences and error variance estimates from multi-centre analysis data,” 2010: M. Wei, Z. Toth, Y. Zhu, Aust. Met. and Ocean Journal.

Dec 2011 – WGNE presentation by Tom Hamill


Analyses from NCEP, ECWMF, UKMO, CMC, FNMOC

00UTC: 1Feb 2008 to 30Apr 2008

7

Wei et al. (2010)

Time-averaged spread over the average anomaly



In general, smaller analysis spread in locations with in-situ observations (esp. radiosondes, aircraft)


Analyses from NCEP, CMC, FNMOC

00UTC, 12UTC: 1Jan 2007 to 1Jun 2007


8

Langland et al. (2008)

500mb Temperature Analyses Root Mean Square Difference (CMC / AVN)



Radiosonde observation count



Indication that assimilation of high-quality in-situ observations (radiosondes, aircraft data) reduces analysis uncertainty more than assimilation of satellite observations (radiances and feature-track or scatterometer winds)


500mb Temperature Analyses

Root Mean Square Difference


1 Jan – 1 Jun 2007

Langland et al. (2008)

NOGAPS / AVN



NOGAPS / CMC




Smaller analysis uncertainty (<1K) where radiosonde data are provided
Larger uncertainty (1-2K) between analyses where satellite data predominates

**UNCERTAINTY BETWEEN ANALYSES CAN BE LARGER THAN SHORT-RANGE
“FORECAST ERROR” !!**

2011: same pattern still in place!

[Many new radiance data have been added during 2007-2011]

Root-Mean Square of Analysis Differences: 500mb Temperature



Analyses from NCEP, ECWMF, UKMO, CMC, CMA


00UTC: 1OCT 2010 to 30Sep 2011

11

Hamill (WGNE, Dec 2011)


500 hPa height

Yearly avg. geopotential height spread (m) at 500 hPa from global analyses



250 hPa u-wind

Yearly avg. u-component spread (m) at 250 hPa from global analyses



Time-average of daily spread (sample standard deviation) of analyses about their daily mean

**“Analyses, assumed to be unbiased, do exhibit substantial bias
Implications for ensemble perturbations (may be too small)”**

300mb Wind Speed (2010) GFS / ECMWF

Root-Mean Square of Analysis Differences: 300mb Wind Speed

2010


GFS | ECMWF

January – December

Langland and Maue 2011

2010 -- 300 hPa Wind Speed Initial Condition Uncertainty


Note the very significant effect of in-situ wind observations:
Radiosondes and Commercial Aircraft



GFS | ECMWF
201

January – September

2011



Radiosonde stations on the budget chopping block

Example: Eareckson Air Station (Shemya) 70414


13

Langland and Maue 2011

Shemya Island




GFS | ECMWF Root mean square difference of analyzed
300mb wind speed: July 2009 – June 2011




Uncertainty in atmospheric upper-tropospheric wind analyses is substantially lower in locations where radiosonde data is provided. The blue-shaded areas are locations where raobs provide soundings twice-daily (00z and 12z). Station 70414 provides data only at 12z, so the associated reduction in analysis error at that location is mitigated, but still significant.


Mean Z500 variance ECMWF NOGAPS METFRANCE GEOS5




Mean Z500 variance ECWMF from average of 4 analyses




Mean Z500 variance MFRANCE from average of 4 analyses



Mean Z500 variance NOGAPS from average of 4 analyses



Mean Z500 variance GEOS5 from average of 4 analyses



Question

Why is analysis uncertainty over oceanic regions still much larger than over North America and Europe, despite the addition of massive amounts of radiance data? [Now as much as 90% of all assimilated data.]

Basic patterns of analysis differences and analysis uncertainty in 2012 remain similar to those reported in 2002

Summary


Assimilation of radiosonde and aircraft data substantially reduces uncertainty in upper-air analyses of temperature and wind

Analysis uncertainty is larger where analyses relies primarily on radiance observations


What new observing instruments and variables are most-needed to reduce analysis uncertainty?

Where is the greatest need to reduce the current magnitude of analysis uncertainty? Polar regions? Oceanic storm tracks? Targeted improvements to observing system and data assimilation procedures?


Mean u250 variance ECMWF NOGAPS METFRANCE GEOS5




Mean u250 variance ECWMF from average of 4 analyses




Mean u250 variance MFRANCE from average of 4 analyses



Mean u250 variance NOGAPS from average of 4 analyses




Mean u250 variance GEOS5 from average of 4 analyses



About 19 million observations assimilated in global domain each day in NAVDAS-AR [4d-Var]



28 Apr 2012 [00, 06, 12, 18 UTC]



HIGH OBSERVATION DENSITY DOES NOT GAURANTEE ANALYSIS QUALITY !!

Count of observations assimilated by NAVDAS-AR

28 Apr 2012 [00, 06, 12, 18 UTC]



Data count in 2° x 2° lat/lon bins

The largest density of observations is due to in-situ data [radiosondes, aircraft, land-surface and ocean-surface observations]